

Lecture 5

Jan 29, 14

- Review and complete Section 1.5
- Section 1.6
 - Valid Arguments and Rules of Inference

Section 1.6 Summary

- Valid Arguments
- Inference Rules for Propositional Logic
- Using Rules of Inference to Build Arguments
- Rules of Inference for Quantified Statements
- Building Arguments for Quantified Statements

Revisiting the Socrates Example

- We have the two premises:
 - “All men are mortal.”
 - “Socrates is a man.”
- And the conclusion:
 - “Socrates is mortal.”
- How do we get the conclusion from the premises?

The Argument

- We can express the premises (above the line) and the conclusion (below the line) in predicate logic as an argument:

$$\forall x (Man(x) \rightarrow Mortal(x))$$

$$Man(Socrates)$$

$$\therefore Mortal(Socrates)$$

- We will see shortly that this is a valid argument.

Valid Arguments

- We will show how to construct valid arguments in two stages; first for propositional logic and then for predicate logic. The rules of inference are the essential building block in the construction of valid arguments.
 1. Propositional Logic
 - Inference Rules
 2. Predicate Logic
 - Inference rules for propositional logic plus additional inference rules to handle variables and quantifiers.

Arguments in Propositional Logic

- A *argument* in propositional logic is a sequence of propositions. All but the final proposition are called *premises*. The last statement is the *conclusion*.
- The argument is valid if the premises imply the conclusion. An *argument form* is an argument that is valid no matter what propositions are substituted into its propositional variables.
- If the premises are $p_1, p_2, \dots, p_n$ and the conclusion is q then $(p_1 \wedge p_2 \wedge \dots \wedge p_n) \rightarrow q$ is a tautology.
- Inference rules are all argument simple argument forms that will be used to construct more complex argument forms.

Rules of Inference for Propositional Logic

Modus Ponens

$$\frac{p \rightarrow q \quad p}{\therefore q}$$

Corresponding Tautology:
 $(p \wedge (p \rightarrow q)) \rightarrow q$

Example:

Let p be “It is snowing.”

Let q be “I will study discrete math.”

“If it is snowing, then I will study discrete math.”

“It is snowing.”

“Therefore, I will study discrete math.”

Modus Tollens

$$\frac{p \rightarrow q \quad \neg q}{\therefore \neg p}$$

Corresponding Tautology:
 $(\neg p \wedge (p \rightarrow q)) \rightarrow \neg q$

Example:

Let p be “it is snowing.”

Let q be “I will study discrete math.”

“If it is snowing, then I will study discrete math.”

“I will not study discrete math.”

“Therefore, it is not snowing.”

Hypothetical Syllogism

$$\frac{p \rightarrow q \quad q \rightarrow r}{\therefore p \rightarrow r}$$

Corresponding Tautology:
 $((p \rightarrow q) \wedge (q \rightarrow r)) \rightarrow (p \rightarrow r)$

Example:

Let p be “it snows.”

Let q be “I will study discrete math.”

Let r be “I will get an A.”

“If it snows, then I will study discrete math.”

“If I study discrete math, I will get an A.”

“Therefore , If it snows, I will get an A.”

Disjunctive Syllogism

$$\begin{array}{c} p \vee q \\ \neg p \\ \hline \therefore q \end{array}$$

Corresponding Tautology:
 $(\neg p \wedge (p \vee q)) \rightarrow q$

Example:

Let p be “I will study discrete math.”

Let q be “I will study English literature.”

“I will study discrete math or I will study English literature.”

“I will not study discrete math.”

“Therefore, I will study English literature.”

Addition

$$\frac{p}{\therefore p \vee q}$$

Corresponding Tautology:
 $p \rightarrow (p \vee q)$

Example:

Let p be “I will study discrete math.”

Let q be “I will visit Las Vegas.”

“I will study discrete math.”

“Therefore, I will study discrete math or I will visit Las Vegas.”

Simplification

$$\frac{p \wedge q}{\therefore q}$$

Corresponding Tautology:
 $(p \wedge q) \rightarrow p$

Example:

Let p be “I will study discrete math.”

Let q be “I will study English literature.”

“I will study discrete math and English literature”

“Therefore, I will study discrete math.”

Conjunction

$$\frac{p}{q} \quad \frac{q}{\therefore p \wedge q}$$

Corresponding Tautology:
 $((p) \wedge (q)) \rightarrow (p \wedge q)$

Example:

Let p be “I will study discrete math.”

Let q be “I will study English literature.”

“I will study discrete math.”

“I will study English literature.”

“Therefore, I will study discrete math and I will study English literature.”

Resolution

Resolution plays an important role in AI and is used in Prolog.

$$\frac{\neg p \vee r \quad p \vee q}{\therefore q \vee r}$$

Corresponding Tautology:
 $((\neg p \vee r) \wedge (p \vee q)) \rightarrow (q \vee r)$

Example:

Let p be “I will study discrete math.”

Let r be “I will study English literature.”

Let q be “I will study databases.”

“I will not study discrete math or I will study English literature.”

“I will study discrete math or I will study databases.”

“Therefore, I will study databases or I will English literature.”

Using the Rules of Inference to Build Valid Arguments

- A *valid argument* is a sequence of statements. Each statement is either a premise or follows from previous statements by rules of inference. The last statement is called conclusion.
- A valid argument takes the following form:

$$\begin{array}{c} S_1 \\ S_2 \\ \cdot \\ \cdot \\ \cdot \\ S_n \\ \therefore C \end{array}$$

Valid Arguments

Example 1: From the single proposition

$$p \wedge (p \rightarrow q)$$

Show that q is a conclusion.

Solution:

Step	Reason
1. $p \wedge (p \rightarrow q)$	Premise
2. p	Conjunction using (1)
3. $p \rightarrow q$	Conjunction using (1)
4. q	Modus Ponens using (2) and (3)

Valid Arguments

Example 2:

- With these hypotheses:
 - “It is not sunny this afternoon and it is colder than yesterday.”
 - “We will go swimming only if it is sunny.”
 - “If we do not go swimming, then we will take a canoe trip.”
 - “If we take a canoe trip, then we will be home by sunset.”
- Using the inference rules, construct a valid argument for the conclusion:
 - “We will be home by sunset.”

Solution:

1. Choose propositional variables:
 - p : “It is sunny this afternoon.” r : “We will go swimming.” t : “We will be home by sunset.”
 - q : “It is colder than yesterday.” s : “We will take a canoe trip.”
2. Translation into propositional logic:

Hypotheses: $\neg p \wedge q, r \rightarrow p, \neg r \rightarrow s, s \rightarrow t$

Conclusion: t

Continued on next slide →

Hypotheses: $\neg p \wedge q, r \rightarrow p, \neg r \rightarrow s, s \rightarrow t$

Conclusion: t

Valid Arguments

3. Construct the Valid Argument

Step	Reason
1. $\neg p \wedge q$	Premise
2. $\neg p$	Simplification using (1)
3. $r \rightarrow p$	Premise
4. $\neg r$	Modus tollens using (2) and (3)
5. $\neg r \rightarrow s$	Premise
6. s	Modus ponens using (4) and (5)
7. $s \rightarrow t$	Premise
8. t	Modus ponens using (6) and (7)

Propositional inference rules table

TABLE 1 Rules of Inference.		
Rule of Inference	Tautology	Name
$\frac{p \quad p \rightarrow q}{\therefore q}$	$(p \wedge (p \rightarrow q)) \rightarrow q$	Modus ponens
$\frac{\neg q \quad p \rightarrow q}{\therefore \neg p}$	$(\neg q \wedge (p \rightarrow q)) \rightarrow \neg p$	Modus tollens
$\frac{p \rightarrow q \quad q \rightarrow r}{\therefore p \rightarrow r}$	$((p \rightarrow q) \wedge (q \rightarrow r)) \rightarrow (p \rightarrow r)$	Hypothetical syllogism
$\frac{p \vee q \quad \neg p}{\therefore q}$	$((p \vee q) \wedge \neg p) \rightarrow q$	Disjunctive syllogism
$\frac{p}{\therefore p \vee q}$	$p \rightarrow (p \vee q)$	Addition
$\frac{p \wedge q}{\therefore p}$	$(p \wedge q) \rightarrow p$	Simplification
$\frac{p \quad q}{\therefore p \wedge q}$	$((p) \wedge (q)) \rightarrow (p \wedge q)$	Conjunction
$\frac{p \vee q \quad \neg p \vee r}{\therefore q \vee r}$	$((p \vee q) \wedge (\neg p \vee r)) \rightarrow (q \vee r)$	Resolution

Problem 3, page 78

3. What rule of inference is used in each of these arguments?

- a) Alice is a mathematics major. Therefore, Alice is either a mathematics major or a computer science major.
- b) Jerry is a mathematics major and a computer science major. Therefore, Jerry is a mathematics major.
- c) If it is rainy, then the pool will be closed. It is rainy. Therefore, the pool is closed.
- d) If it snows today, the university will close. The university is not closed today. Therefore, it did not snow today.
- e) If I go swimming, then I will stay in the sun too long. If I stay in the sun too long, then I will sunburn. Therefore, if I go swimming, then I will sunburn.

Choices:

TABLE 1 Rules of Inference.		
Rule of Inference	Tautology	Name
$\frac{p \quad p \rightarrow q}{\therefore q}$	$(p \wedge (p \rightarrow q)) \rightarrow q$	Modus ponens
$\frac{\neg q \quad p \rightarrow q}{\therefore \neg p}$	$(\neg q \wedge (p \rightarrow q)) \rightarrow \neg p$	Modus tollens
$\frac{p \rightarrow q \quad q \rightarrow r}{\therefore p \rightarrow r}$	$((p \rightarrow q) \wedge (q \rightarrow r)) \rightarrow (p \rightarrow r)$	Hypothetical syllogism
$\frac{p \vee q \quad \neg p}{\therefore q}$	$((p \vee q) \wedge \neg p) \rightarrow q$	Disjunctive syllogism
$\frac{p}{\therefore p \vee q}$	$p \rightarrow (p \vee q)$	Addition
$\frac{p \wedge q}{\therefore p}$	$(p \wedge q) \rightarrow p$	Simplification
$\frac{p \quad q}{\therefore p \wedge q}$	$((p) \wedge (q)) \rightarrow (p \wedge q)$	Conjunction
$\frac{p \vee q \quad \neg p \vee r}{\therefore q \vee r}$	$((p \vee q) \wedge (\neg p \vee r)) \rightarrow (q \vee r)$	Resolution

Problem 3, page 78

3. What rule of inference is used in each of these arguments?

- a) Alice is a mathematics major. Therefore, Alice is either a mathematics major or a computer science major.
- b) Jerry is a mathematics major and a computer science major. Therefore, Jerry is a mathematics major.
- c) If it is rainy, then the pool will be closed. It is rainy. Therefore, the pool is closed.
- d) If it snows today, the university will close. The university is not closed today. Therefore, it did not snow today.
- e) If I go swimming, then I will stay in the sun too long. If I stay in the sun too long, then I will sunburn. Therefore, if I go swimming, then I will sunburn.

- a) Addition
- b) Simplification
- c) Modus ponens
- d) Modus tollens
- e) Hypothetical syllogism

Choices:

TABLE 1 Rules of Inference.		
Rule of Inference	Tautology	Name
$\frac{p \quad p \rightarrow q}{\therefore q}$	$(p \wedge (p \rightarrow q)) \rightarrow q$	Modus ponens
$\frac{\neg q \quad p \rightarrow q}{\therefore \neg p}$	$(\neg q \wedge (p \rightarrow q)) \rightarrow \neg p$	Modus tollens
$\frac{p \rightarrow q \quad q \rightarrow r}{\therefore p \rightarrow r}$	$((p \rightarrow q) \wedge (q \rightarrow r)) \rightarrow (p \rightarrow r)$	Hypothetical syllogism
$\frac{p \vee q \quad \neg p}{\therefore q}$	$((p \vee q) \wedge \neg p) \rightarrow q$	Disjunctive syllogism
$\frac{p}{\therefore p \vee q}$	$p \rightarrow (p \vee q)$	Addition
$\frac{p \wedge q}{\therefore p}$	$(p \wedge q) \rightarrow p$	Simplification
$\frac{p \quad q}{\therefore p \wedge q}$	$((p) \wedge (q)) \rightarrow (p \wedge q)$	Conjunction
$\frac{p \vee q \quad \neg p \vee r}{\therefore q \vee r}$	$((p \vee q) \wedge (\neg p \vee r)) \rightarrow (q \vee r)$	Resolution

Handling Quantified Statements

- Valid arguments for quantified statements are a sequence of statements. Each statement is either a premise or follows from previous statements by rules of inference which include:
 - Rules of Inference for Propositional Logic
 - Rules of Inference for Quantified Statements
- The rules of inference for quantified statements are introduced in the next several slides.

Universal Instantiation (UI)

$$\frac{\forall x P(x)}{\therefore P(c)}$$

Example:

Our domain consists of all dogs and Fido is a dog.

“All dogs are cuddly.”

“Therefore, Fido is cuddly.”

Universal Generalization (UG)

$$\frac{P(c) \text{ for an arbitrary } c}{\therefore \forall x P(x)}$$

Used often implicitly in Mathematical Proofs.

Existential Instantiation (EI)

$$\frac{\exists x P(x)}{\therefore P(c) \text{ for some element } c}$$

Example:

“There is someone who got an A in the course.”

“Let’s call her a and say that a got an A”

Existential Generalization (EG)

$$\frac{P(c) \text{ for some element } c}{\therefore \exists x P(x)}$$

Example:

“Michelle got an A in the class.”

“Therefore, someone got an A in the class.”

Using Rules of Inference

Example 1: Using the rules of inference, construct a valid argument to show that

“John Smith has two legs”

is a consequence of the premises:

“Every man has two legs.” “John Smith is a man.”

Solution: Let $M(x)$ denote “ x is a man” and $L(x)$ “ x has two legs” and let John Smith be a member of the domain.

Valid Argument:

Step	Reason
1. $\forall x(M(x) \rightarrow L(x))$	Premise
2. $M(J) \rightarrow L(J)$	UI from (1)
3. $M(J)$	Premise
4. $L(J)$	Modus Ponens using (2) and (3)

Using Rules of Inference

Example 2: Use the rules of inference to construct a valid argument showing that the conclusion

“Someone who passed the first exam has not read the book.”
follows from the premises

“A student in this class has not read the book.”

“Everyone in this class passed the first exam.”

Solution: Let $C(x)$ denote “ x is in this class,” $B(x)$ denote “ x has read the book,” and $P(x)$ denote “ x passed the first exam.”

First we translate the
premises and conclusion
into symbolic form.

$$\frac{\begin{array}{l} \exists x(C(x) \wedge \neg B(x)) \\ \forall x(C(x) \rightarrow P(x)) \end{array}}{\therefore \exists x(P(x) \wedge \neg B(x))}$$

Continued on next slide →

Using Rules of Inference

Valid Argument:

Step	Reason
1. $\exists x(C(x) \wedge \neg B(x))$	Premise
2. $C(a) \wedge \neg B(a)$	EI from (1)
3. $C(a)$	Simplification from (2)
4. $\forall x(C(x) \rightarrow P(x))$	Premise
5. $C(a) \rightarrow P(a)$	UI from (4)
6. $P(a)$	MP from (3) and (5)
7. $\neg B(a)$	Simplification from (2)
8. $P(a) \wedge \neg B(a)$	Conj from (6) and (7)
9. $\exists x(P(x) \wedge \neg B(x))$	EG from (8)

Returning to the Socrates Example

$$\forall x (Man(x) \rightarrow Mortal(x))$$

$$Man(Socrates)$$

$$\therefore Mortal(Socrates)$$

Solution for Socrates Example

Valid Argument

Step

1. $\forall x(Man(x) \rightarrow Mortal(x))$
2. $Man(Socrates) \rightarrow Mortal(Socrates)$
3. $Man(Socrates)$
4. $Mortal(Socrates)$

Reason

Premise
UI from (1)
Premise
MP from (2)
and (3)

Universal Modus Ponens

Universal Modus Ponens combines universal instantiation and modus ponens into one rule.

$$\forall x(P(x) \rightarrow Q(x))$$

$P(a)$, where a is a particular
element in the domain

$$\therefore Q(a)$$

This rule could be used in the Socrates example.