

# Math 141

## Lecture 13

Greg Maloney

Todor Milev

University of Massachusetts Boston

Spring 2015

# Outline

## 1 Alternating Series

- Estimating Sums
- Absolute Convergence

## 2 Absolute Convergence and the Ratio and Root Tests

- The Ratio Test
- The Root Test

# License to use and redistribute

These lecture slides and their  $\text{\LaTeX}$  source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project (a notice of use freecalc is sufficient).

- Latest version of the .tex sources of the slides: `https://sourceforge.net/p/freecalculus/code/HEAD/tree/`
- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:  
`https://creativecommons.org/licenses/by/3.0/us/`  
and the links therein.

# Alternating Series

## Definition (Alternating Series)

An alternating series is a series whose terms are alternately positive and negative.

## Examples

Here are two examples:

$$1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \frac{1}{5} - \frac{1}{6} + \cdots = \sum_{n=1}^{\infty} (-1)^{n-1} \frac{1}{n}$$

$$-\frac{1}{2} + \frac{2}{3} - \frac{3}{4} + \frac{4}{5} - \frac{5}{6} + \frac{6}{7} - \cdots = \sum_{n=1}^{\infty} (-1)^n \frac{n}{n+1}$$

The  $n$ th term of an alternating series has the form

$$a_n = (-1)^{n-1} b_n \quad \text{or} \quad a_n = (-1)^n b_n$$

where  $b_n$  is positive.

## Theorem (The Alternating Series Test)

*If the alternating series*

$$\sum_{n=1}^{\infty} (-1)^{n-1} b_n = b_1 - b_2 + b_3 - b_4 + b_5 - \cdots, \quad b_n > 0$$

*satisfies*

- ①  $b_{n+1} \leq b_n$  for all  $n$  and
- ②  $\lim_{n \rightarrow \infty} b_n = 0$

*then the series is convergent.*

## Example

The alternating harmonic series

$$1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \cdots = \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n}$$

satisfies

①  $b_{n+1} < b_n$  because  $\frac{1}{n+1} < \frac{1}{n}$ .

②  $\lim_{n \rightarrow \infty} b_n = \lim_{n \rightarrow \infty} \frac{1}{n} = 0$ .

Therefore the series is convergent by the Alternating Series Test.

## Example

The series  $\sum_{n=1}^{\infty} (-1)^n \frac{3n}{4n-1}$  is alternating, but

$$\lim_{n \rightarrow \infty} b_n = \lim_{n \rightarrow \infty} \frac{3n}{4n-1} \cdot \frac{1}{\frac{1}{n}} = \lim_{n \rightarrow \infty} \frac{3}{4 - \frac{1}{n}} = \frac{3}{4}$$

Therefore the series is divergent by the Alternating Series Test.

# Estimating Sums

This theorem allows us to estimate the size of the remainder  $R_n = s - s_n$  in an alternating series.

## Theorem (Alternating Series Estimation Theorem)

Let  $\sum (-1)^{n-1} b_n$  be the sum of an alternating series that satisfies

①  $0 \leq b_{n+1} \leq b_n$  and

②  $\lim_{n \rightarrow \infty} b_n = 0$ .

Then the size of the error is less than the first omitted term; that is,

$$|R_n| = |s - s_n| \leq b_{n+1}.$$



## Example

Find the sum of  $\sum_{n=0}^{\infty} \frac{(-1)^n}{n!}$  correct to three decimal places. ( $0! = 1$ .)

$$\textcircled{1} \quad b_{n+1} = \frac{1}{(n+1)!} = \frac{1}{n!(n+1)} < \frac{1}{n!} = b_n.$$

$$\textcircled{2} \quad 0 < \frac{1}{n!} < \frac{1}{n} \rightarrow 0, \text{ so } b_n \rightarrow 0 \text{ as } n \rightarrow \infty.$$

- Therefore the series converges by the Alternating Series Test.

$$\begin{aligned} s &= \frac{1}{0!} - \frac{1}{1!} + \frac{1}{2!} - \frac{1}{3!} + \frac{1}{4!} - \frac{1}{5!} + \frac{1}{6!} - \frac{1}{7!} + \cdots \\ &= 1 - 1 + \frac{1}{2} - \frac{1}{6} + \frac{1}{24} - \frac{1}{120} + \frac{1}{720} - \frac{1}{5040} + \cdots \end{aligned}$$

- $|s - s_6| \leq b_7 = \frac{1}{5040} < 0.0002.$
- $s_6 = 1 - 1 + \frac{1}{2} - \frac{1}{6} + \frac{1}{24} - \frac{1}{120} + \frac{1}{720} \approx 0.368056.$
- The error of less than 0.0002 doesn't affect the third decimal place, so  $s \approx s_6 \approx 0.368.$

# Absolute Convergence and the Ratio and Root Tests

In this section, we start with any series  $\sum a_n$  and consider the corresponding series

$$\sum |a_n| = |a_1| + |a_2| + |a_3| + \cdots$$

consisting of the absolute values of the terms of the original series.

# Absolute Convergence

## Definition (Absolutely Convergent)

A series  $\sum a_n$  is called absolutely convergent if the series of absolute values  $\sum |a_n|$  is convergent.

If  $\sum a_n$  is a series with all positive terms, then  $|a_n| = a_n$  and absolute convergence is the same thing as convergence in this case.

## Example

The series

$$\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n^2} = 1 - \frac{1}{2^2} + \frac{1}{3^2} - \frac{1}{4^2} + \cdots$$

is absolutely convergent because

$$\sum_{n=1}^{\infty} \left| \frac{(-1)^{n-1}}{n^2} \right| = 1 + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \cdots$$

is a convergent  $p$ -series with  $p = 2$ .

## Example

The alternating harmonic series

$$\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} = 1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \cdots$$

is convergent (by the alternating series test, as already demonstrated).

- Is it absolutely convergent?

$$\sum_{n=1}^{\infty} \left| \frac{(-1)^{n-1}}{n} \right| = 1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \cdots$$

- This is a  $p$ -series with  $p = 1$ .
- Therefore  $\sum_{n=1}^{\infty} \left| \frac{(-1)^{n-1}}{n} \right|$  is divergent.
- Therefore  $\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n}$  is not absolutely convergent.

## Definition (Conditionally Convergent)

A series  $\sum a_n$  is called conditionally convergent if it is convergent but not absolutely convergent.

- The alternating harmonic series is conditionally convergent.
- Therefore it is possible for a series to be convergent but not absolutely convergent.
- Question: Is it possible for a series to be absolutely convergent but not convergent?
- Answer: No. This is the content of the next theorem.

## Theorem (Absolute Convergence Implies Convergence)

*If a series is absolutely convergent, then it is convergent.*

## Example

Determine whether

$$\sum_{n=1}^{\infty} \frac{\cos n}{n^2} = \frac{\cos 1}{1^2} + \frac{\cos 2}{2^2} + \frac{\cos 3}{3^2} + \frac{\cos 4}{4^2} + \cdots$$

is convergent or divergent.

- The series has positive and negative terms, but is not alternating.
- Use the Comparison Test:

$$\begin{array}{rcl} 0 & \leq & |\cos n| \leq 1 \\ 0 & \leq & \frac{|\cos n|}{n^2} \leq \frac{1}{n^2} \end{array}$$

- $\sum \frac{1}{n^2}$  is a  $p$ -series with  $p = 2$ .
- Therefore  $\sum \frac{1}{n^2}$  is convergent, and so by the Comparison Test,  $\sum \frac{|\cos n|}{n^2}$  is also convergent.
- Therefore  $\sum \frac{\cos n}{n^2}$  is absolutely convergent.
- Therefore by the previous theorem,  $\sum \frac{\cos n}{n^2}$  is convergent.



# The Ratio Test

## Theorem (The Ratio Test)

- 1 If  $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = L < 1$ , then the series  $\sum a_n$  is absolutely convergent (and therefore convergent).
- 2 If  $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = L > 1$  or  $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \infty$ , then the series  $\sum a_n$  is divergent.
- 3 If  $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = L = 1$ , then the Ratio Test is inconclusive.

The Ratio Test is inconclusive if  $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = 1$ .

## Example

$$\sum_{n=1}^{\infty} \frac{1}{n^2}$$

- This is a  $p$ -series with  $p = 2$ .
- Therefore it is convergent.

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{\frac{1}{(n+1)^2}}{\frac{1}{n^2}} = \frac{n^2}{(n+1)^2} \cdot \frac{1}{n^2} = \frac{1}{\left(1 + \frac{1}{n}\right)^2} \rightarrow 1 \quad \text{as } n \rightarrow \infty$$

## Example

$$\sum_{n=1}^{\infty} \frac{1}{n}$$

- This is a  $p$ -series with  $p = 1$ .
- Therefore it is divergent.

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{\frac{1}{n+1}}{\frac{1}{n}} = \frac{n}{n+1} \cdot \frac{1}{n} = \frac{1}{1 + \frac{1}{n}} \rightarrow 1 \quad \text{as } n \rightarrow \infty$$

## Example

Test the series  $\sum_{n=1}^{\infty} (-1)^n \frac{n^3}{3^n}$  for absolute convergence.

$$\begin{aligned} \left| \frac{a_{n+1}}{a_n} \right| &= \left| \frac{(-1)^{n+1} \frac{(n+1)^3}{3^{n+1}}}{(-1)^n \frac{n^3}{3^n}} \right| \\ &= \frac{(n+1)^3}{3^{n+1}} \cdot \frac{3^n}{n^3} \\ &= \frac{1}{3} \left( \frac{n+1}{n} \right)^3 \\ &= \frac{1}{3} \left( 1 + \frac{1}{n} \right)^3 \\ &\rightarrow \frac{1}{3} < 1 \end{aligned}$$

Therefore the series is absolutely convergent by the Ratio Test.

## Example

Test the convergence of the series  $\sum_{n=1}^{\infty} \frac{n^n}{n!}$ .

$$\begin{aligned} \left| \frac{a_{n+1}}{a_n} \right| &= \left| \frac{\frac{(n+1)^{n+1}}{(n+1)!}}{\frac{n^n}{n!}} \right| \\ &= \frac{(n+1)^{n+1}}{(n+1)!} \cdot \frac{n!}{n^n} \\ &= \frac{(n+1)(n+1)^n}{(n+1)n!} \cdot \frac{n!}{n^n} \\ &= \left( \frac{n+1}{n} \right)^n = \left( 1 + \frac{1}{n} \right)^n \\ &\rightarrow e > 1 \end{aligned}$$

Therefore the series is divergent by the Ratio Test.

# The Root Test

## Theorem (The Root Test)

- 1 If  $\lim_{n \rightarrow \infty} \sqrt[n]{|a_n|} = L < 1$ , then the series  $\sum a_n$  is absolutely convergent (and therefore convergent).
- 2 If  $\lim_{n \rightarrow \infty} \sqrt[n]{|a_n|} = L > 1$  or  $\lim_{n \rightarrow \infty} \sqrt[n]{|a_n|} = \infty$ , then the series  $\sum a_n$  is divergent.
- 3 If  $\lim_{n \rightarrow \infty} \sqrt[n]{|a_n|} = L = 1$ , then the Root Test is inconclusive.

If  $L = 1$  in the Ratio Test, don't try the Root Test, because it will be inconclusive too.

If  $L = 1$  in the Root Test, don't try the Ratio Test, because it will be inconclusive too.

## Example

Test convergence of the series  $\sum_{n=1}^{\infty} \left( \frac{2n+3}{3n+2} \right)^n$ .

$$a_n = \left( \frac{2n+3}{3n+2} \right)^n$$

$$\sqrt[n]{|a_n|} = \frac{2n+3}{3n+2} \cdot \frac{1}{n}$$

$$= \frac{2 + \frac{3}{n}}{3 + \frac{2}{n}}$$

$$\rightarrow \frac{2}{3} < 1$$

Therefore the series is absolutely convergent by the Root Test.