



University of Connecticut
Department of Mathematics

MATH 2210

CHAPTER 1 WORKSHEET

SPRING 2019

Due: Thursday, March 7

Directions: This worksheet covers Chapter 1 from Lay's Linear Algebra book. Turn in the red-squared problems below (problems 3, 5, 6, 10, 11, 12, 15, 20, 21, and 23) on separate paper. Use only one side of the paper, remove all fringes, and staple this cover sheet to your submission.

*Any of these problems is fair game for the exam (not just the red ones).
Ask questions if you're unsure about anything.*

GROUP MEMBER(S) (up to four)

NAME: _____

NAME: _____

NAME: _____

NAME: _____

Section (of the student I will hand this back to): _____


Checklist for formatting:

- ☐ write every group member's name
- ☐ remove all fringes
- ☐ use only one side of the paper
- ☐ label each problem clearly
- ☐ give a clear justification for your answer
- ☐ circle your answer
- ☐ staple this coversheet to your submission

CHAPTER 1


1. Mark each statement True or False. Justify each answer. (If true, cite appropriate facts or theorems. If false, explain why or give a counterexample that shows why the statement is not true in every case.)
 - a. Every matrix is row equivalent to a unique matrix in echelon form.
 - b. Any system of n linear equations in n variables has at most n solutions.
 - c. If a system of linear equations has two different solutions, it must have infinitely many solutions.
 - d. If a system of linear equations has no free variables, then it has a unique solution.
 - e. If an augmented matrix $[A \ \mathbf{b}]$ is transformed into $[C \ \mathbf{d}]$ by elementary row operations, then the equations $A\mathbf{x} = \mathbf{b}$ and $C\mathbf{x} = \mathbf{d}$ have exactly the same solution sets.
 - f. If a system $A\mathbf{x} = \mathbf{b}$ has more than one solution, then so does the system $A\mathbf{x} = \mathbf{0}$.
 - g. If A is an $m \times n$ matrix and the equation $A\mathbf{x} = \mathbf{b}$ is consistent for some $\mathbf{b}$, then the columns of A span $\mathbb{R}^m$.
 - h. If an augmented matrix $[A \ \mathbf{b}]$ can be transformed by elementary row operations into reduced echelon form, then the equation $A\mathbf{x} = \mathbf{b}$ is consistent.
 - i. If matrices A and B are row equivalent, they have the same reduced echelon form.
 - j. The equation $A\mathbf{x} = \mathbf{0}$ has the trivial solution if and only if there are no free variables.
 - k. If A is an $m \times n$ matrix and the equation $A\mathbf{x} = \mathbf{b}$ is consistent for every $\mathbf{b}$ in $\mathbb{R}^m$, then A has m pivot columns.
 - l. If an $m \times n$ matrix A has a pivot position in every row, then the equation $A\mathbf{x} = \mathbf{b}$ has a unique solution for each $\mathbf{b}$ in $\mathbb{R}^m$.
 - m. If an $n \times n$ matrix A has n pivot positions, then the reduced echelon form of A is the $n \times n$ identity matrix.
 - n. If 3×3 matrices A and B each have three pivot positions, then A can be transformed into B by elementary row operations.
 - o. If A is an $m \times n$ matrix, if the equation $A\mathbf{x} = \mathbf{b}$ has at least two different solutions, and if the equation $A\mathbf{x} = \mathbf{c}$ is consistent, then the equation $A\mathbf{x} = \mathbf{c}$ has many solutions.
 - p. If A and B are row equivalent $m \times n$ matrices and if the columns of A span $\mathbb{R}^m$, then so do the columns of B .
 - q. If none of the vectors in the set $S = \{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ in $\mathbb{R}^3$ is a multiple of one of the other vectors, then S is linearly independent.
 - r. If $\{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$ is linearly independent, then $\mathbf{u}$, $\mathbf{v}$, and $\mathbf{w}$ are not in $\mathbb{R}^2$.
 - s. In some cases, it is possible for four vectors to span $\mathbb{R}^5$.
 - t. If $\mathbf{u}$ and $\mathbf{v}$ are in $\mathbb{R}^m$, then $-\mathbf{u}$ is in $\text{Span}\{\mathbf{u}, \mathbf{v}\}$.
 - u. If $\mathbf{u}$, $\mathbf{v}$, and $\mathbf{w}$ are nonzero vectors in $\mathbb{R}^2$, then $\mathbf{w}$ is a linear combination of $\mathbf{u}$ and $\mathbf{v}$.
 - v. If $\mathbf{w}$ is a linear combination of $\mathbf{u}$ and $\mathbf{v}$ in $\mathbb{R}^n$, then $\mathbf{u}$ is a linear combination of $\mathbf{v}$ and $\mathbf{w}$.
 - w. Suppose that $\mathbf{v}_1, \mathbf{v}_2$, and $\mathbf{v}_3$ are in $\mathbb{R}^5$, $\mathbf{v}_2$ is not a multiple of $\mathbf{v}_1$, and $\mathbf{v}_3$ is not a linear combination of $\mathbf{v}_1$ and $\mathbf{v}_2$. Then $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ is linearly independent.
 - x. A linear transformation is a function.
 - y. If A is a 6×5 matrix, the linear transformation $\mathbf{x} \mapsto A\mathbf{x}$ cannot map $\mathbb{R}^5$ onto $\mathbb{R}^6$.
 - z. If A is an $m \times n$ matrix with m pivot columns, then the linear transformation $\mathbf{x} \mapsto A\mathbf{x}$ is a one-to-one mapping.
2. Let a and b represent real numbers. Describe the possible solution sets of the (linear) equation $ax = b$. [Hint: The number of solutions depends upon a and b .]
3. The solutions (x, y, z) of a single linear equation $ax + by + cz = d$ form a plane in $\mathbb{R}^3$ when a, b , and c are not all zero. Construct sets of three linear equations whose graphs (a) intersect in a single line, (b) intersect in a single point, and (c) have no

points in common. Typical graphs are illustrated in the figure.




Three planes intersecting in a line

(a)




Three planes intersecting in a point

(b)



Three planes with no intersection

(c)



Three planes with no intersection

(c')

4. Suppose the coefficient matrix of a linear system of three equations in three variables has a pivot position in each column. Explain why the system has a unique solution.

5. Determine h and k such that the solution set of the system (i) is empty, (ii) contains a unique solution, and (iii) contains infinitely many solutions.

$$\begin{array}{ll} \text{a. } x_1 + 3x_2 = k & \text{b. } -2x_1 + hx_2 = 1 \\ 4x_1 + hx_2 = 8 & 6x_1 + kx_2 = -2 \end{array}$$

6. Consider the problem of determining whether the following system of equations is consistent:

$$\begin{array}{l} 4x_1 - 2x_2 + 7x_3 = -5 \\ 8x_1 - 3x_2 + 10x_3 = -3 \end{array}$$

- Define appropriate vectors, and restate the problem in terms of linear combinations. Then solve that problem.
 - Define an appropriate matrix, and restate the problem using the phrase “columns of A .”
 - Define an appropriate linear transformation T using the matrix in (b), and restate the problem in terms of T .
7. Consider the problem of determining whether the following system of equations is consistent for all b_1, b_2, b_3 :

$$\begin{array}{l} 2x_1 - 4x_2 - 2x_3 = b_1 \\ -5x_1 + x_2 + x_3 = b_2 \\ 7x_1 - 5x_2 - 3x_3 = b_3 \end{array}$$

- Define appropriate vectors, and restate the problem in terms of $\text{Span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$. Then solve that problem.
- Define an appropriate matrix, and restate the problem using the phrase “columns of A .”


- Define an appropriate linear transformation T using the matrix in (b), and restate the problem in terms of T .

8. Describe the possible echelon forms of the matrix A . Use the notation of Example 1 in Section 1.2.

- A is a 2×3 matrix whose columns span $\mathbb{R}^2$.
- A is a 3×3 matrix whose columns span $\mathbb{R}^3$.

9. Write the vector $\begin{bmatrix} 5 \\ 6 \end{bmatrix}$ as the sum of two vectors, one on the line $\{(x, y) : y = 2x\}$ and one on the line $\{(x, y) : y = x/2\}$.

10. Let $\mathbf{a}_1, \mathbf{a}_2$, and $\mathbf{b}$ be the vectors in $\mathbb{R}^2$ shown in the figure, and let $A = [\mathbf{a}_1 \ \mathbf{a}_2]$. Does the equation $A\mathbf{x} = \mathbf{b}$ have a solution? If so, is the solution unique? Explain.



11. Construct a 2×3 matrix A , not in echelon form, such that the solution of $A\mathbf{x} = \mathbf{0}$ is a line in $\mathbb{R}^3$.

12. Construct a 2×3 matrix A , not in echelon form, such that the solution of $A\mathbf{x} = \mathbf{0}$ is a plane in $\mathbb{R}^3$.

13. Write the *reduced* echelon form of a 3×3 matrix A such that the first two columns of A are pivot columns and
- $$A \begin{bmatrix} 3 \\ -2 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}.$$

14. Determine the value(s) of a such that $\left\{ \begin{bmatrix} 1 \\ a \end{bmatrix}, \begin{bmatrix} a \\ a+2 \end{bmatrix} \right\}$ is linearly independent.

15. In (a) and (b), suppose the vectors are linearly independent. What can you say about the numbers $a, \dots, f$? Justify your answers. [Hint: Use a theorem for (b).]

$$\text{a. } \begin{bmatrix} a \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} b \\ c \\ 0 \end{bmatrix}, \begin{bmatrix} d \\ e \\ f \end{bmatrix} \quad \text{b. } \begin{bmatrix} a \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} b \\ c \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} d \\ e \\ f \\ 1 \end{bmatrix}$$

16. Use Theorem 7 in Section 1.7 to explain why the columns of the matrix A are linearly independent.

$$A = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 2 & 5 & 0 & 0 \\ 3 & 6 & 8 & 0 \\ 4 & 7 & 9 & 10 \end{bmatrix}$$

17. Explain why a set $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{v}_4\}$ in $\mathbb{R}^5$ must be linearly independent when $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ is linearly independent and $\mathbf{v}_4$ is *not* in $\text{Span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$.

18. Suppose $\{\mathbf{v}_1, \mathbf{v}_2\}$ is a linearly independent set in $\mathbb{R}^n$. Show that $\{\mathbf{v}_1, \mathbf{v}_1 + \mathbf{v}_2\}$ is also linearly independent.

19. Suppose $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$ are distinct points on one line in $\mathbb{R}^3$. The line need not pass through the origin. Show that $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ is linearly dependent.

20. Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a linear transformation, and suppose $T(\mathbf{u}) = \mathbf{v}$. Show that $T(-\mathbf{u}) = -\mathbf{v}$.


21. Let $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be the linear transformation that reflects each vector through the plane $x_2 = 0$. That is, $T(x_1, x_2, x_3) = (x_1, -x_2, x_3)$. Find the standard matrix of T .

22. Let A be a 3×3 matrix with the property that the linear transformation $\mathbf{x} \mapsto A\mathbf{x}$ maps $\mathbb{R}^3$ onto $\mathbb{R}^3$. Explain why the transformation must be one-to-one.

23. A *Givens rotation* is a linear transformation from $\mathbb{R}^n$ to $\mathbb{R}^n$ used in computer programs to create a zero entry in a vector (usually a column of a matrix). The standard matrix of a Givens rotation in $\mathbb{R}^2$ has the form

$$\begin{bmatrix} a & -b \\ b & a \end{bmatrix}, \quad a^2 + b^2 = 1$$

Find a and b such that $\begin{bmatrix} 4 \\ 3 \end{bmatrix}$ is rotated into $\begin{bmatrix} 5 \\ 0 \end{bmatrix}$.



A Givens rotation in $\mathbb{R}^2$.

24. The following equation describes a Givens rotation in $\mathbb{R}^3$. Find a and b .

$$\begin{bmatrix} a & 0 & -b \\ 0 & 1 & 0 \\ b & 0 & a \end{bmatrix} \begin{bmatrix} 2 \\ 3 \\ 4 \end{bmatrix} = \begin{bmatrix} 2\sqrt{5} \\ 3 \\ 0 \end{bmatrix}, \quad a^2 + b^2 = 1$$