

## Localized electric field manipulates a nuclear spin

Rachel Berkowitz

Citation: [Physics Today](#) **73**, 5, 16 (2020); doi: 10.1063/PT.3.4468

View online: <https://doi.org/10.1063/PT.3.4468>

View Table of Contents: <https://physicstoday.scitation.org/toc/pto/73/5>

Published by the [American Institute of Physics](#)

---

### ARTICLES YOU MAY BE INTERESTED IN

[An atomic physics perspective on the kilogram's new definition](#)

[Physics Today](#) **73**, 32 (2020); <https://doi.org/10.1063/PT.3.4472>

[Radar points the way to detecting cosmic neutrinos](#)

[Physics Today](#) **73**, 14 (2020); <https://doi.org/10.1063/PT.3.4467>

[A single-atom heat engine](#)

[Physics Today](#) **73**, 66 (2020); <https://doi.org/10.1063/PT.3.4482>

[Tying celestial mechanics to Earth's ice ages](#)

[Physics Today](#) **73**, 48 (2020); <https://doi.org/10.1063/PT.3.4474>

[Murray Gell-Mann](#)

[Physics Today](#) **73**, 63 (2020); <https://doi.org/10.1063/PT.3.4480>

[A physicist out of academia](#)

[Physics Today](#) **73**, 12 (2020); <https://doi.org/10.1063/PT.3.4466>

---



## Solve for the Unknown

**APSIT**  
INSURANCE FOR SCIENCE PROFESSIONALS

**SEE THE SOLUTION**

look for ionization trails from neutrino-induced cascades. In nature, those cascades take place inside the ice. But in the proof-of-concept experiment, the researchers had to address the transition radiation. Fortunately, that RF noise was similar from pulse to pulse.

To extract persuasive evidence of a cascade reflection, the researchers filtered out of their data the transition radiation and other noise—Askaryan RF fields, telecommunication signals, and reflections from concrete and metal features in the SLAC station, shown in figure 2. They performed three types of experiments: ones with both the electron beam and radar on; ones with the radar

on but not the electron beam; and ones with the electron beam on but not the radar. Armed with those data, they subtracted the background to resolve a real radar signal. To constrain the analysis, they confirmed that the signal had the expected timing, frequency, and power dependence.

Prohira and his colleagues next want to repeat the experiment on a high-altitude ice sheet in Antarctica. It's radio quiet there—though even the passage of wind generates residual RF hum—and the altitude increases the likelihood that a cosmic-ray-induced cascade will make it into the ice; the ionization trail will come from that cascade. Antennas just

below the surface would transmit radar and pick up reflected signals.

After that in-nature test the researchers will turn their attention to neutrino-induced cascades.

Mark Wilson

## References


1. S. Prohira et al., *Phys. Rev. Lett.* **124**, 091101 (2020).
2. G. A. Askaryan, *Sov. Phys. JETP* **14**, 441 (1962).
3. P. M. S. Blackett, A. C. B. Lovell, *Proc. R. Soc. London A* **177**, 183 (1941).
4. K. D. de Vries, K. Hanson, T. Meures, *Astropart. Phys.* **60**, 25 (2015).
5. S. Prohira et al., *Phys. Rev. D* **100**, 072003 (2019).

# Localized electric field manipulates a nuclear spin

The atom-level control could provide the precision required for some quantum computing applications.

**O**ne promising approach for quantum information processing involves embedding tightly spaced arrays of identical atomic nuclei in a silicon substrate. In that design, each nucleus's spin serves as a quantum bit, or qubit. The qubit's spin, which can be set to different states, is used to store and process information. However, before spin-based devices can be scaled up for practical use, quantum engineers need to be able to control a single nuclear spin in silicon without affecting adjacent spins.

In principle, NMR could do the job. Radio-frequency (RF) magnetic field pulses can excite and control nuclear spins that are polarized in a static magnetic field. Because of the pulses' wide spatial extent, however, they tend to influence adjacent spins, which renders NMR impractical for manipulating individual spins in a collection of identical atoms. Ideally, a method for controlling individual nuclear spins would match the ease of exciting individual electron spins in a row of semiconductor quantum dots, in which each dot is equipped with a separate electrode. Adapting that approach for nuclear spins offers a potential advantage because nuclear spins




have longer coherence times than electron spins and can be measured with minimal readout error.

Electric fields, rather than magnetic ones, provide an intriguing possibility for nuclear spin control. The fields can be efficiently routed and tightly confined in complex nanoscale devices. Indeed, highly focused electric fields make possible the sophisticated interconnections found in modern silicon computer chips.

Now researchers in Andrea Morello's

**FIGURE 1. IN THIS ARTIST'S IMPRESSION** of a nuclear electric resonance device, a sharp metallic antenna applies a strong oscillating electric field directly to an antimony atom (green) embedded in a silicon chip. Other metallic components include electrostatic gate connections and readout electrodes. (Image by Tony Melov/UNSW.)

lab at the University of New South Wales in Australia have demonstrated electrical control over nuclear spin.<sup>1</sup> The re-



**FIGURE 2. A SCANNING ELECTRON MICROGRAPH** shows the metal-oxide-semiconductor device used in the experiment. An antimony donor atom is implanted in the silicon substrate. Electrostatic gates control the potential of the Sb atom embedded in the chip between the gates. The device contains a single-electron transistor (SET) for spin readout and a microwave antenna carrying an electric current that normally generates an oscillating magnetic field for performing NMR. (Adapted from ref. 1.)

searchers developed a technique for controlling the spin of a single atom embedded in a silicon chip, illustrated in figure 1, by using an electric field produced at the tip of a nanometer-sized electrode.

## Bulk efforts

The suggestion that electric fields could control nuclear spins dates to the 1960s.<sup>2</sup> Nico Bloembergen predicted that an external electric field can induce spins larger than  $\frac{1}{2}$  to transition from one energy level to another in crystals with low symmetry. Such nuclei can have nonspherical charge density distributions, or quadrupole moments, which couple to local electric field gradients. When placed in a crystalline environment without inversion symmetry, a time-varying external electric field should modulate that coupling and change the spin state.

Bloembergen's efforts to demonstrate electrical control over spin saw partial success. His studies proved that static electric fields could shift spins between energy levels in a bulk crystal.<sup>3</sup>

"heroic experiments," says Morello, the investigations were unable to create the requisite huge RF electric fields in bulk crystals.

Only recently did experiments clearly demonstrate nuclear electric resonance (NER) in a bulk crystal. In 2013 Masaaki Ono and colleagues at Tohoku University, Japan, excited spins in gallium arsenide at their resonant frequency using an oscillating electric field.<sup>4</sup> Ono's team proved the viability of using NER to manipulate spins in a bulk crystal.


## Electrical accident

Coherent control over a single nuclear spin in a silicon device remained an unsolved problem. To investigate possible solutions for single spin control, Morello's PhD student Serwan Asaad and postdoc Vincent Mourik used an antenna to magnetically probe an antimony-123 nucleus embedded in silicon. The researchers had originally set out to perform NMR on a single  $^{123}\text{Sb}$  nucleus, which has spin  $\frac{1}{2}$ . They fabricated the de-

vice, shown in figure 2, with an Sb atom and a special antenna optimized to create a high-frequency magnetic field to control the nucleus. But when they applied power to the antenna, they observed a resonance spectrum that was missing several expected NMR peaks.

Asaad realized that the high power had damaged the antenna and transformed it into an open circuit. As a result, the antenna was creating a strong electric field focused directly on the Sb atom. Conveniently, the device also included several tiny electrodes with which to control the potential of the atom. The researchers redesigned their experiments to apply an oscillating voltage to one of the electrodes and observed the same resonance spectrum as when applying power to the damaged antenna. Then, they calculated the rate at which transitions between different energy-dependent spin states should occur in response to either a magnetic or an electric field (see figure 3). Comparing those calculations to experimental data persuaded them that the electric field was causing the Sb resonances.

Still, the result puzzled the researchers. The electric quadrupole moment—the nonspherical charge distribution of the nucleus—does not couple directly to an electric field, but to the field's gradient. To understand the effect, Morello enlisted Andrew Baczewski of Sandia National Laboratories to perform *ab initio* calculations of the phenomenon. The simulations revealed that the applied electric field distorted the charge distributions in the bonds between the Sb atom and the Si atoms around it. Then



**FIGURE 3. AN RF FIELD** drives antimony-123 atom transitions between different nuclear spin states at different rates, the so-called Rabi frequency. The measured rates (colored circles) match theoretical predictions for transitions due to an electric field (NER, stars) but not those due to a magnetic field (NMR, triangles). (Adapted from ref. 1.)



the distortion of the charges produced a strongly nonuniform electric field around the Sb nucleus. That field provided the gradient that caused the nuclear spin transitions. The calculated effects match the experimental results.

Because electric fields decay rapidly with distance from the electrode and can easily be screened with other metallic structures, NER allows control that is localized enough to manipulate individual

nuclei. The technique could help scale up a key component in quantum computing applications, according to Anthony Sigillito from Princeton University. The level of localized control demonstrated by Morello and coworkers could provide the precision needed to drive one qubit without affecting the other in a two-qubit gate, which is the fundamental building block of a scalable quantum computer. The new result opens the pos-

sibility of all-electrical driving of a donor qubit embedded in a silicon device.

Rachel Berkowitz

## References

1. S. Asaad et al., *Nature* **579**, 205 (2020).
2. *Science* **133**, 1363 (1961), abstract by N. Bloembergen.
3. R. W. Dixon, N. Bloembergen, *J. Chem. Phys.* **41**, 1739 (1964).
4. M. Ono et al., *Appl. Phys. Express* **6**, 033002 (2013).

# Spongy hydrogels clean textured paintings

Washing away the dirt from an artistic masterpiece is especially tricky when the surface is not flat. New materials can help.

There will never be another Vincent van Gogh, Pablo Picasso, Georgia O'Keeffe, or Frida Kahlo. Each of the great works of visual art that together make up humankind's cultural heritage is unique and irreplaceable. Conservators at museums thus face a pair of often contradictory tasks: Keep the artwork in their care safe for future generations and allow as many people as possible to enjoy it now.

Even under the best of conditions, a painting on display for decades accumulates dirt and dust that mar its appearance and dull its colors. Although cleaning everyday grime from ordinary objects isn't technically challenging—scrubbing with soap and water usually does the trick—priceless works of art require a gentler, more sophisticated touch to avoid any risk of damage. Worse, many artists, especially in recent decades, apply paint to canvas in thick brush strokes to create three-dimensional textured paintings, like the one in figure 1a, with lots of nooks and crannies where dirt can hide.

Now Piero Baglioni and colleagues at the University of Florence in Italy have developed a polymer hydrogel—a net-

**FIGURE 1. TEXTURAL FEATURES** of contemporary paintings can be bumpy, (a) as shown by the uncovered mock-up. (b) Most cleaning gels, including this one developed for restoring paper artwork,<sup>2</sup> are unable to cover the rough surfaces uniformly. (c) A gel of polyvinyl alcohol, however, is soft enough to do the job. (Images adapted from ref. 1.)

work of polymer chains bound together into a porous, water-bearing solid—that safely removes dirt from the roughest of painted surfaces.<sup>1</sup> “These gels are unique in the conservation field,” says Bronwyn Ormsby, the principal conservation scientist at Tate, a network of art museums in the UK. “They can offer solutions to some of our more intractable problems.”

## Freeze-thaw gels

Pastes, poultices, thickeners, and absorbent materials have been used for thousands of years to apply fluids to surfaces. You'd probably clean up a mess using a wet rag or sponge rather than dousing it with soapy water. Confining a liquid in a solid or semisolid matrix prevents it from immediately inundating the surface and flowing away.

The use of gels for cleaning art and artifacts, which dates back to the 1980s, is based on a similar principle. A sheet of gel, imbued with water, surfactant solution, or other solvent, is gently placed onto the surface to be cleaned and left there for a minute or two. During that



time, tiny amounts of fluid seep out of the gel and loosen the dirt particles. The gel is then peeled away, and with it, one hopes, comes the dirt.

Conventional gels—many of which



## From Tight Spaces to Tight Tolerances

we machine sapphire,  
ceramics, quartz & glass

www.insaco.com | 215.536.3500



Earning your trust since 1947